arXiv:1011.6011v2 [math.DS] 17 Dec 2010

Hyperbolic Measures with Dominated Splitting

Xueting Tian

Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing
100190, China

E-mail: *tianxt@amss.ac.cn*

Abstract

In this paper we consider a non-atomic invariant hyperbolic measure μ of a C^1 diffeomorphsim on a compact manifold, in whose Oseledec splitting the stable bundle dominates the unstable bundle on μ a.e. points. We show an *exponentially* shadowing and an *exponentially* closing lemma, and as applications we show two classical results. One is that there exists a hyperbolic periodic point such that the closure of its unstable manifold has *positive* measure and it has a homoclinic point from which one can deduce a horseshoe. Moreover, such hyperbolic periodic points are dense in the support $supp(\mu)$ of the given hyperbolic measure. Another is to show Livshitz Theorem.

1 Introduction

Let M be a compact D-dimensional smooth Riemannian manifold and let ρ denote the distance induced by the Riemannian metric. Denote the tangent bundle of M by TM. Denote by $\mathrm{Diff}^1(M)$ the space of C^1 diffeomorphisms of M. Hereafter let $f \in \mathrm{Diff}^1(M)$. Before stating our result we firstly introduce some notions.

We begin with the notions of invariant measure and ergodic measure. A Borel measure μ is said to be $f-$invariant, if $\mu(f^{-1}(B)) = \mu(B)$ for all measurable sets B. An $f-$invariant measure μ is said to be f ergodic, if the only measurable sets B with $f^{-1}(B) = B$ satisfies $\mu(B) = 0$ or $\mu(B) = 1$. Given an invariant measure μ, by the Oseledec theorem [12], there is a Borel set $L(\mu)$ satisfying $fL(\mu) = L(\mu)$ and $\mu(L(\mu)) = 1$, called Oseledec basin of μ, such that for every $x \in L(\mu)$, there exist
(a) real numbers, called Lyapunov exponents,

$$\lambda_1(x) < \lambda_2(x) < \cdots < \lambda_{W(x)}(x) \ (W(x) \leq D) \tag{1.1}$$

(b) positive integers $m_1(x), \ \cdots, \ \ m_{W(x)}(x)$, satisfying $m_1(x) + \cdots + m_{W(x)}(x) = D$;
(c) a measurable splitting $T_xM = E^1_x \oplus \cdots \oplus E^{W(x)}_x$ with $\dim E^i_x = m_i(x)$ and $Df(E^i_x) = E^i_{f(x)}$, such that

$$\lim_{n\to\pm\infty} \frac{\log \|Df^n v\|}{n} = \lambda_i(x),$$

Key words and phrases: Lyapunov exponents, Pesin set, Liao's closing lemma, stable manifold, C^1 non-uniform hyperbolicity, horseshoe, dominated splitting, periodic point and periodic measure

AMS Review: 37D25, 37D30, 37D10, 37C50, 37C25.

with uniform convergence on $\{v \in E_x^i \mid \|v\| = 1\},\ i = 1,\, 2,\, \cdots,\, W(x)$.
In particular, if μ is ergodic, $\lambda_i(x), m_i(x),\ W(x)$ can always be constants for μ a.e. x.

Now we recall the notion of hyperbolic invariant measure.

Definition 1.1. *We call an f invariant measure μ to be hyperbolic, if*
(1) none of the Lyapunov exponents for μ are zero;
(2) there exist Lyapunov exponents with different signs.

If μ is an f hyperbolic invariant measure, then from Oseledec theorem above there is $R(x) \in \mathbb{N}$ such that $\lambda_{R(x)} < 0 < \lambda_{R(x)+1}$ in the inequality (1.1). Let

$$E^s(x) := E_x^1 \oplus \cdots \oplus E_x^{R(x)}$$

and

$$E^u(x) := E_x^{R(x)+1} \oplus \cdots \oplus E_x^{W(x)},$$

then we get a measurable $Df-$invariant splitting $T_xM = E^s(x) \oplus E^u(x)$ on $L(\mu)$, called *Oseledec's hyperbolic splitting* of μ. We call $E^s(\cdot)$ to be the stable bundle and call $E^u(\cdot)$ to be the unstable bundle of μ.

Definition 1.2. *We call an f invariant measure μ to be hyperbolic with a fixed index, if μ is hyperbolic and for μ a.e. x, the dimension of the stable bundle(composed by Oseledec bundles whose Lyapunov exponents are negative) are constant, i.e., $dim(E^s(x)) \equiv i$ for some positive integer i.*

Denote the minimal norm of an invertible linear map A by $m(A) = \|A^{-1}\|^{-1}$. We recall the notion of dominated splitting. Let Δ be an $f-$invariant set and $T_\Delta M = E \oplus F$ be a $Df-$invariant splitting on Δ. $T_\Delta M = E \oplus F$ is called to be (S_0, λ)-dominated on Δ (or simply dominated), if there exist two constants $S_0 \in \mathbb{Z}^+$ and $\lambda > 0$ such that

$$\frac{1}{S} \log \frac{\|Df^S|_{E(x)}\|}{m(Df^S|_{F(x)})} \le -2\lambda,\ \forall x \in \Delta,\ S \ge S_0.$$

Note that dominated splitting can be extended to its closure, even neighborhoods and dominated splitting is always continuous (see [2]). Let $x \in M$ and $T_{Orb(x)}M = E \oplus F$ be a $Df-$invariant splitting on the orbit $Orb(x)$ of x. We call $T_{Orb(x)}M = E \oplus F$ to be dominated at x, if $T_{Orb(x)}M = E \oplus F$ is (S_0, λ)-dominated on the set $\Delta = Orb(x)$ for some $S_0 = S_0(x) \in \mathbb{Z}^+$ and $\lambda = \lambda(x) > 0$. Note that dominated splittings at different points admit different S_0 and λ.

To state (exponentially) shadowing lemma and closing lemma we need some notions. Given $x \in M$ and $n \in \mathbb{N}$, let

$$\{x,\, n\} := \{f^j(x) \mid j = 0,\, 1,\, \cdots,\, n\}.$$

In other words, $\{x,\, n\}$ represents the orbit segment from x to $f^n(x)$ with length n. For a sequence of points $\{x_i\}_{i=-\infty}^{+\infty}$ in M and a sequence of positive integers $\{n_i\}_{i=-\infty}^{+\infty}$, we call $\{x_i,\, n_i\}_{i=-\infty}^{+\infty}$ a δ-pseudo-orbit, if $\rho(f^{n_i}(x_i),\, x_{i+1}) < \delta$ for all i. Given $\theta > 0$ and $\eta > 0$, we call a point $x \in M$ an (exponentially) (η, θ)-shadowing point for a pseudo-orbit $\{x_i,\, n_i\}_{i=-\infty}^{+\infty}$, if

$$\rho\big(f^{c_i+j}(x), f^j(x_i)\big) < \eta \cdot e^{-\min\{j, n_i-j\}\theta},$$

$\forall\ j = 0,\, 1,\, 2,\, \cdots,\, n_i$ and $\forall\ i \in \mathbb{Z}$, where c_i is defined as

$$c_i = \begin{cases} 0, & \text{for } i = 0 \\ \sum_{j=0}^{i-1} n_j, & \text{for } i > 0 \\ -\sum_{j=i}^{-1} n_j, & \text{for } i < 0. \end{cases} \tag{1.2}$$

We call a point $x \in M$ an η-shadowing point for a pseudo-orbit $\{x_i,\, n_i\}_{i=-\infty}^{+\infty}$, if

$$\rho\big(f^{c_i+j}(x), f^j(x_i)\big) < \eta,$$

$\forall\ j = 0,\, 1,\, 2,\, \cdots,\, n_i$ and $\forall\ i \in \mathbb{Z}$.

Since hyperbolic invariant measure can be always divided into at most $D = dim(M)$ hyperbolic invariant measures with fixed indexs, for convenience, in this section we always suppose that

"$f \in \mathrm{Diff}^1(M)$ and preserves a nonatomic invariant hyperbolic measure μ with a *fixed index* & the Oseledec's hyperbolic splitting $T_xM = E^s(x) \oplus E^u(x)$ is dominated at μ a.e. x."

Now we start to state our main results as follows.

Theorem 1.3. *For each $\tau > 0$, there exist a compact set $\Lambda_\tau \subseteq M$, $\theta_\tau > 0$ and $T_\tau \in \mathbb{N}$ such that $\mu(\Lambda_\tau) > 1 - \tau$ and following two properties hold.*
(i) (Exponentially) Shadowing Lemma*: For $\forall\ \eta > 0$, there exists $\delta = \delta(\tau, \eta) > 0$ such that if a δ-pseudo-orbit $\{x_i,\, n_i\}_{i=-\infty}^{+\infty}$ satisfies $n_i \ge T_\tau$ and $x_i, f^{n_i}(x_i) \in \Lambda_\tau$ for all i, then there exists an (exponentially) (η, θ_τ)-shadowing point $x \in M$ for $\{x_i, n_i\}_{i=-\infty}^{+\infty}$. If further $\{x_i, n_i\}_{i=-\infty}^{+\infty}$ is periodic, i.e., there exists an integer $m > 0$ such that $x_{i+m} = x_i$ and $n_{i+m} = n_i$ for all i, then the shadowing point x can be chosen to be periodic.*
(ii) Stable Manifold Theorem*: There exists $\sigma > 0$ such that all points in $\Lambda_\tau^*(\sigma)$ have uniform sizes of stable and unstable manifolds, where $\Lambda_\tau^*(\sigma)$ denotes the union set of Λ_τ and the set*

$$\{z \in M \mid \exists \textit{ pseudo-orbit } \{x_i,\, n_i\}_{i=-\infty}^{+\infty} \textit{ satisfying } n_i \ge T_\tau \textit{ and } x_i, f^{n_i}(x_i) \in \Lambda_\tau$$

$$\textit{for all } i \textit{ such that } z \textit{ is a } \sigma - \textit{shadowing point for } \{x_i, n_i\}_{i=-\infty}^{+\infty}\}.$$

Moreover, if $x, y \in \Lambda_\tau^*(\sigma)$ and are close enough, then the (local) stable manifold x is transverse to the (local) unstable manifold of y.

Remark 1.4. The assumption of fixed index is to provide that the dimensions of the stable bundles are always consistent. In the present paper we always assume that the dimensions of the stable bundles are all the same.

Theorem 1.5. *(Exponentially Closing lemma) For each $\tau > 0$, there exist a compact set $\Lambda_\tau \subseteq M$, $\theta_\tau > 0$ and $T_\tau \in \mathbb{N}$ such that $\mu(\Lambda_\tau) > 1 - \tau$ and following two properties hold.*
(i) (Exponentially) closing Lemma*: For $\forall\ \eta > 0$, there exists $\beta = \beta(\tau, \eta) > 0$ such that if for an orbit segment $\{x, n\}$ with length $n \geq T_\tau$, one has $x, f^n(x) \in \Lambda_\tau$ and $\rho(x, f^n(x)) < \beta$, then there exists a hyperbolic periodic point $z = z(x) \in M$ satisfying:*

(1) $f^n(z) = z$;

(2) $\rho(f^j(x),\ f^j(z)) < \eta \cdot e^{-\min\{j, n-j\}\theta_\tau}$, $j = 0, 1, \cdots, n-1$.

(ii) Stable Manifold Theorem*: There exists $\sigma > 0$ such that all points in $\Lambda_\delta^*(\sigma)$ have uniform sizes of stable and unstable manifolds, where $\Lambda_\tau^*(\sigma)$ denotes the union set of Λ_τ and the set of periodic points nearby Λ_τ*

$$\{z \in M \mid \exists\, x, n \geq T_\tau \text{ with } x, f^n(x) \in \Lambda_\tau \text{ s.t. } f^n(z) = z,\ \rho(f^i(x),\ f^i(z)) \leq \sigma, 0 \leq i \leq n\}.$$

Moreover, if $x, y \in \Lambda_\tau^*(\sigma)$ are close enough, then the (local) stable manifold x is transverse to the (local) unstable manifold of y.

As consequences of Theorem 1.5, we show two results. One is about the positive-measured property of the closure of unstable manifolds of hyperbolic periodic points and another is about Livshitz Theorem.

Theorem 1.6. *There exists a hyperbolic periodic point with homoclinic point (Hence there exists horseshoe) such that the closure of its global unstable manifold has positive measure. Moreover, the support of the measure μ is contained in the closure of such hyperbolic periodic points.*

Remark 1.7. In the $C^{1+\alpha}$ case, the existence of horseshoes can be found in [13, 7] and the result that the closure of unstable manifold of the shadowing periodic point is of positive measure can be found in [16] by Ugarocovici. Moreover, we point out that we can also use closing lemma to get the second result in [16], since closing lemma is enough to prove(Here we delete the details). Some discussion of existences of periodic points with homoclinic points for *ergodic* hyperbolic measures with dominated splitting of C^1 diffeomorphisms also appeared in [5, 18].

In fact, we can find that $\eta-$closing(i.e., $\rho(f^i(x),\ f^i(z)) < \eta$, $i = 0, 1, \cdots, n-1$) is enough to the proof of Theorem 1.6. However, the proofs of some other properties need the inequality (2) of Theorem 1.5. For example, we show such a theorem(Livshitz Theorem) as follows, which is obtained by Katok [8] for the $C^{1+\alpha}$ case (also see [10, 3]).

Theorem 1.8. *Let $\varphi : M \to \mathbb{R}$ be a Höder continuous function such that for each (hyperbolic) periodic point z with $f^m(z) = z$, we have $\sum_{i=0}^{m-1} \varphi(f^i(z)) = 0$. Then there exists a Borel measurable function ψ such that*

$$\varphi(x) = \psi(f(x)) - \psi(x)$$

holds for μ almost every point x.

This paper is organized as follows. In Section 2 and 3 we introduce stable manifold theorem in the C^1 case and in Section 4 we use this stable manifold theorem and the shadowing lemma in [15] to get exponentially shadowing lemma. In Section 5 we give a proof of Theorem 1.3 and in Section 6 we give the proof of Theorem 1.6 and 1.8.

2 Dominated Splitting and (Center) Stable Manifold

Let $I_1 = (-1,1)$ be the open unit ball and $I_\epsilon = \{x \in \mathbb{R} : |x| < \epsilon\}$ the open $\epsilon-$ball. Denote by $Emb(I_1, M)$ the set of C^1 embeddings of I_1 in M, equipped with the (uniform) C^1 topology. In this section we always suppose that
" Δ is a closed $f-$invariant set & there is a dominated splitting $T_\Delta M = E \oplus F$ over Δ."

The following lemma is taken from [6] about the existence of center stable and unstable manifolds for a dominated splitting.

Lemma 2.1. *([6]) There exist two continuous functions* $\theta^{cs} : \Delta \to Emb(I_1, M)$ *and* $\theta^{cu} : \Delta \to Emb(I_1, M)$ *such that the following properties hold:*
(a) $\theta^{cs}(x)(0) = x$ *and* $\theta^{cu}(x)(0) = x$.
(b) $T_x W^{cs}_\epsilon(x) = E(x)$ *and* $T_x W^{cu}_\epsilon(x) = F(x)$, *where* $W^{cs}_\epsilon(x) = \theta^{cs}(x)I_\epsilon$ *and* $W^{cu}_\epsilon(x) = \theta^{cu}(x)I_\epsilon$.
(c) For $0 < \epsilon_1 < 1$ *there exists* $0 < \epsilon_2 < 1$ *such that*

$$f(W^{cs}_{\epsilon_2}(x)) \subseteq W^{cs}_{\epsilon_1}(f(x))$$

and

$$f^{-1}(W^{cu}_{\epsilon_2}(x)) \subseteq W^{cu}_{\epsilon_1}(f^{-1}(x))$$

for all $x \in \Delta$.

The manifold W^{cs} is called the (local) center stable manifold and W^{cu} the (local) center unstable manifold. They are only locally invariant. Generally, this kind of invariant manifold is not unique. So when we use center invariant manifolds for analysis, we will (henceforth) fix a family in discs in Lemma 2.1.

Now let us consider whenever the center stable and unstable manifold are the truth ones.

Proposition 2.2. *For* $K \in \mathbb{N}$, $C > 0$ *and* $0 < \lambda < \bar{\lambda} < 1$, *there exist* $\epsilon_0 = \epsilon(K, C, \lambda, \bar{\lambda}) > 0$ *and* $\bar{C} = \bar{C}(K, C, \bar{\lambda}) \geq 1$ *such that if for* $x \in \Delta$, *there is a two-sided sequence of integers*

$$\cdots < t_{-2} < t_{-1} < t_0 = 0 < t_1 < t_2 < \cdots$$

with $t_i - t_{i-1} \leq K$ *for all* i *satisfying*

$$\prod_{j=0}^{i-1} \|Df^{t_{j+1}-t_j}|_{E(f^{t_j}(x))}\| \leq C\lambda^{t_i}$$

and

$$\prod_{j=0}^{-i+1} \|Df^{t_{j-1}-t_j}|_{F(f^{t_j}(x))}\| \leq C\lambda^{-t_{-i}} (\textit{in other words, } \prod_{j=-i}^{-1} m(Df^{t_{j+1}-t_j}|_{F(f^{t_j}(x))}) \geq C\lambda^{t_{-i}})$$

for all $i \in \mathbb{K}$, *then for any* $0 < \epsilon \leq \epsilon_0$

$$diam(f^n W^{cs}_\epsilon(x)) \to_n 0 \qquad and \qquad diam(f^{-n} W^{cu}_\epsilon(x)) \to_n 0,$$

i.e., the center stable manifold of size ϵ is in fact a stable manifold and the center unstable manifold of size ϵ is in fact an unstable manifold. Moreover, for all $n \geq 1$,

$$\frac{diam(f^n W^{cs}_\epsilon(x))}{diam(W^{cs}_\epsilon(x))} \leq \bar{C}\bar{\lambda}^n$$

and

$$\frac{diam(f^{-n} W^{cu}_\epsilon(x))}{diam(W^{cu}_\epsilon(x))} \leq \bar{C}\bar{\lambda}^n.$$

Proof of Proposition 2.2 Let Δ be a compact invariant set and $T_\Delta M = E \oplus F$ be a $Df-$invariant dominated splitting over Δ. By Lemma 2.1 the center stable and unstable manifolds exist. Here fix a small constant $\tau \in (1, \frac{\bar{\lambda}}{\lambda})$. One chooses a small constant $\delta > 0$ such that for any $x \in \Delta$, any point $y \in W^{cs}_1(x)$ with $\rho(x, y) < \delta$ and any vector $v \in T_y W^{cs}_1(x)$ and $1 \leq j \leq L$ we have

$$\|D_y f^j \cdot v\| \leq \tau^j \, \|Df^j|_{E(x)}\| \cdot \|v\|.$$

Let $C_f > 1$ be a bound on the norm of the derivative Df. Define

$$\bar{C} = \max\{C_f^K \cdot C \cdot \bar{\lambda}^{-K},\, 1\}.$$

Let $\epsilon_0 > 0$ small enough such that

$$diam(W^{cs/cu}_{\epsilon_0}(y) < \frac{\delta}{\bar{C}}$$

for all $y \in \Delta$. Let $0 < \epsilon \leq \epsilon_0$. Write $n = t_i + q$ where $0 \leq q < t_{i+1} - t_i \leq K$.Since

$$\prod_{j=0}^{i-1} \|Df^{t_{j+1}-t_j}|_{E(f^{t_j}(x))}\| \leq C\lambda^{t_i},$$

then one deduces inductively on n,

$$\begin{aligned} & diam(f^n W^{cs}_\epsilon(x)) \\ \leq\ & C_f^q \cdot \Pi_{j=0}^{i-1} \tau^{(t_{j+1}-t_j)} \, \|Df^{t_{j+1}-t_j}|_{E(f^{t_j}(x))}\| \cdot diam(W^{us}_\epsilon(x)) \\ \leq\ & C_f^K \cdot \tau^{t_i} \cdot C\lambda^{t_i} \cdot diam(W^{us}_\epsilon(x)) \\ \leq\ & \bar{C}\bar{\lambda}^n \cdot diam(W^{cs}_\epsilon(x)). \end{aligned}$$

Note that

$$diam(f^n W^{cs}_\epsilon(x)) \leq \bar{C}\bar{\lambda}^n \cdot \frac{\delta}{\bar{C}} = \bar{\lambda}^n \cdot \delta \leq \delta,$$

so that the induction can go on and on. Moreover,

$$diam(f^n W^{cs}_\epsilon(x)) \leq \bar{\lambda}^n \cdot \delta,$$

which goes to 0 as $n \to +\infty$.

Similarly, one has get the result for the unstable manifold. □

The following corollary is a direct consequence of the continuity of $W^{cs/cu}(x)$.

Corollary 2.3. *There exists* $\epsilon_0 > 0$ *such that for any* $0 < \epsilon \le \epsilon_0$, *there exists* $\delta = \delta(\epsilon)$ *such that for any* $x, y \in \Delta$, *if* $\rho(x, y) < \delta$, *then* $W^{cs}_\epsilon(x)$ *and* $W^{cu}_\epsilon(y)$ *are transversal and have a unique intersection point.*

Furthermore, we also have such a corollary directly from condition (c) in Lemma 2.1 and the continuity of $W^{cs/cu}(x)$.

Corollary 2.4. *There exists* $\epsilon_0 > 0$ *such that for any* $0 < \epsilon \le \epsilon_0$, *there exists* $\delta = \delta(\epsilon)$ *such that for any* $x \in \Delta$,
a) if $y \in W^{cs}_\epsilon(x)$ *and* $\rho(f^{-1}(x), f^{-1}(y)) < \delta$ *then* $f^{-1}(y) \in W^{cu}_\epsilon(f^{-1}(x))$;
b) if $y \in W^{cu}_\epsilon(x)$ *and* $\rho(f(x), f(y)) < \delta$ *then* $f(y) \in W^{cu}_\epsilon(f(x))$.

Remark 2.5. Similar discussions of above results in this section for C^1 *surface* diffeomorphisms also appeared in [14, 5].

3 Stable manifolds on Pesin blocks

In this section all statements are independent on measures. The main aim of this section is preparing stable manifold theorem on Pesin blocks for proving Theorem 1.3.

3.1 Pesin blocks, Pesin set

In this subsection we recall the definitions of Pesin blocks and Pesin set from [15](with little modification).

Definition 3.1. *Given* $K \in \mathbb{N}$, $\zeta > 0$, *and for all* $k \in \mathbb{Z}^+$, *we define* $\Lambda_k = \Lambda_k(K, \zeta)$ *to be all points* $x \in M$ *for which there is a splitting* $T_xM = E(x) \oplus F(x)$ *with the invariance property* $D_xf(E(x)) = E(f(x))$ *and* $D_xf(F(x)) = F(f(x))$ *and satisfying:*

(a).
$$\frac{\log \|Df^r|_{E(x)}\| + \sum_{j=0}^{l-1} \log \|Df^K|_{E(f^{jK+r}(x))}\|}{lK + r} \le -\zeta, \qquad \forall\ l \ge k,\ \forall\ 0 \le r \le K-1;$$

(b).
$$\frac{\log m(Df^r|_{F(f^{-lK-r}(x))}) + \sum_{j=-l}^{-1} \log m(Df^K|_{F(f^{jK}(x))})}{lK + r} \ge \zeta, \qquad \forall\ l \ge k,\ \forall\ 0 \le r \le K-1;$$

(c).
$$\frac{1}{L} \log \frac{\|Df^L|_{E(f^l(x))}\|}{m(Df^L|_{F(f^l(x))})} \le -2\zeta, \quad \forall\ l \in \mathbb{Z},\ L \ge K.$$

Denote by $\Lambda = \Lambda(K, \zeta)$ *the maximal* $f-$*invariant subset of* $\bigcup_{k\geq 1} \Lambda_k$*, meaning*

$$\Lambda = \bigcap_{n\in\mathbb{Z}} f^n(\bigcup_{k\geq 1} \Lambda_k).$$

We call Λ *a Pesin set and call* $\Lambda_k (k \geq 1)$ *Pesin blocks. (see Figure 1 to explain (a) and (b), respectively).*



Figure 1: Graph to show (a) and (b).

Remark 3.2. 1. Every Pesin block in present paper is contained in the original Pesin set defined in [15], since conditions (a) and (b) are same and the condition (c) implies the original condition (c) in [15].

2. Following similar discussion in [15], we also have that $\Lambda_k(K, \zeta)$ are an increasing sequence of closed subsets of M. The fact that the splitting is continuous on $\Lambda_k(K, \zeta)$ is obvious, since the splitting is dominated on the invariant set $\cup_{n\in\mathbb{Z}} f^n \Lambda_k(K, \zeta)$ by condition (c) and dominated splitting is always continuous[2].

Now let us define a set which contains the Pesin block and the points nearby Pesin block. Let $\sigma > 0$. Define $\Lambda^*_k(K, \zeta, \sigma)$ as the union set of $\Lambda_k(K, \zeta)$ and

$$\{z \in M \mid \exists \text{ pseudo-orbit } \{x_i,\, n_i\}_{i=-\infty}^{+\infty} \text{ satisfying } n_i \geq 2kK \text{ and } x_i, f^{n_i}(x_i) \in \Lambda_k(K, \zeta)$$

$$\text{for all } i \text{ such that } z \text{ is a } \sigma-\text{shadowing point for } \{x_i, n_i\}_{i=-\infty}^{+\infty}\}.$$

3.2 Uniform size of Stable Manifolds on Pesin blocks

From condition (c) in the definition of Pesin block Λ_k, the two bundles are dominated on invariant set $\cup_{l\in\mathbb{Z}} f^l(\Lambda_k)$ and then on its closure, since dominated property can always be extended on the closure even neighborhoods. Thus, combining condition (1) and (2) it is easy to get the following corollary from Proposition 2.2, which shows that for every Pesin block Λ_k, the (un)stable manifolds of $x \in \Lambda_k$ exist and have uniform length.

Corollary 3.3. *For $K \in \mathbb{K}$, $\zeta > \bar{\zeta} > 0$ and $k \geq 1$, there exist $\epsilon_0 = \epsilon(K, \zeta, \bar{\zeta}, k) > 0$ and $\bar{C} = \bar{C}(K, k, \zeta, \bar{\zeta}) \geq 1$ such that for any $0 < \epsilon \leq \epsilon_0$ and for all $x \in \Lambda_k = \Lambda_k(K, \zeta)$, $diam(f^n W_\epsilon^{cs}(x)) \to_n 0$ and $diam(f^{-n} W_\epsilon^{cu}(x)) \to_n 0$, i.e., the center stable manifold of size ϵ is in fact a stable manifold and the center unstable manifold of size ϵ is in fact an unstable manifold. Moreover, for all $n \geq 1$,*

$$\frac{diam(f^n W_\epsilon^{cs}(x))}{diam(W_\epsilon^{cs}(x))} \leq \bar{C} e^{-\bar{\zeta} n}$$

and

$$\frac{diam(f^{-n} W_\epsilon^{cu}(x))}{diam(W_\epsilon^{cu}(x))} \leq \bar{C} e^{-\bar{\zeta} n}.$$

Proof of Corollary 3.3 Let $\Delta := \overline{\cup_{l \in \mathbb{Z}} f^l(\Lambda_k)}$. Clearly Δ is a compact invariant set and $T_\Delta M = E \oplus F$ be a $Df-$invariant dominated splitting over Δ, due to condition (c) in the definition of Pesin block Λ_k.

Let $\lambda := \lambda(\zeta) = e^{-\zeta}$, $\bar{\lambda} := \bar{\lambda}(\bar{\zeta}) = e^{-\bar{\zeta}}$ and define $C := C(\zeta, k, K) = \{e^\zeta \cdot C_f\}^{kK}$ where $C_f > 1$ be a bound on the norm of the derivative Df. Then for $x \in \Lambda_k$, by the first and second condition in the definition of Λ_k one has

$$\Pi_{i=0}^{l-1} \| Df^K|_{E(f^{iK}x)} \| \leq C \lambda^{-lK}$$

and

$$\Pi_{i=0}^{l-1} \| Df^{-K}|_{E(f^{-iK}x)} \| \leq C \lambda^{-lK}$$

for all $l \geq 1$. Clearly for all $x \in \Lambda_k$, the sequence in Proposition 2.2 can always be chosen $\{t_i = iK\}_{i=-\infty}^{+\infty}$. So using Proposition 2.2, the stable and unstable manifolds on Λ_k can be chosen of uniform size only dependent on $K, C, \lambda, \bar{\lambda}$ and thus only dependent on $K, \zeta, \bar{\zeta}, k$. So we can complete the proof. $\square$

Moreover, we can get uniform size for stable and unstable manifolds for all points in $\Lambda_k^*(K, \zeta, \sigma)$ if σ is small enough.

Proposition 3.4. *For $K \in \mathbb{K}$, $\zeta > \bar{\zeta} > 0$ and $k \geq 1$, there exist $\sigma = \sigma(K, \zeta, \bar{\zeta}, k) > 0, \epsilon_0 = \epsilon(K, \zeta, \bar{\zeta}, k) > 0$ and $\bar{C} = \bar{C}(K, k, \zeta, \bar{\zeta}) > 0$ such that*
(1) For any $0 < \epsilon \leq \epsilon_0$ and for all $x \in \Lambda_k^*(K, \zeta, \sigma)$, $diam(f^n W_\epsilon^{cs}(x)) \to_n 0$ and $diam(f^{-n} W_\epsilon^{cu}(x)) \to_n 0$, i.e., the center stable manifold of size ϵ is in fact a stable manifold and the center unstable manifold of size ϵ is in fact an unstable manifold. Moreover, for all $n \geq 1$,

$$\frac{diam(f^n W_\epsilon^{cs}(x))}{diam(W_\epsilon^{cs}(x))} \leq \bar{C} e^{-\bar{\zeta} n}$$

and

$$\frac{diam(f^{-n} W_\epsilon^{cu}(x))}{diam(W_\epsilon^{cu}(x))} \leq \bar{C} e^{-\bar{\zeta} n}.$$

(2) For any $0 < \epsilon \leq \epsilon_0$, there exists $\delta = \delta(\epsilon)$ such that for any $x, y \in \Lambda_k^*(K, \zeta, \sigma)$ if $\rho(x, y) < \delta$, then $W_\epsilon^{cs}(x)$ and $W_\epsilon^{cu}(y)$ are transversal and have a unique intersection point, and thus the stable manifold and unstable manifold are transversal and have a intersection point.

Proof Let $\epsilon_0^1 = \epsilon_0(\zeta, \bar{\zeta}, k, K)$ and $\bar{C}_1 = \bar{C}(K, k, \zeta, \bar{\zeta}) > 0$ be the ϵ_0 and $\bar{C}$ as in Corollary 3.3 and thus the center stable and center unstable manifolds of all $x \in \Lambda_k(K, \zeta)$ are the truth stable and unstable manifolds, and their sizes are at least ϵ_0^1.

Let $\Delta_1 := \overline{\cup_{l\in\mathbb{Z}} f^l(\Lambda_k)}$. Clearly Δ_1 is a compact invariant set and $T_{\Delta_1}M = E \oplus F$ be a $Df-$invariant dominated splitting over Δ_1, due to condition (c) in the definition of Pesin block Λ_k. Since dominated property can be extended to its neighborhoods (see [2]), we can take an open neighborhood $U = B_{\Delta_1}(\tau) = \{x \mid \rho(x, \Delta_1) < \tau\}$ (for some small $\tau > 0$) of Δ_1 such that the splitting is extended and dominated on $\Delta := \cap_{n\in\mathbb{Z}} f^n(U)$. Take and fix $\zeta > \hat{\zeta} > \bar{\zeta}$. Let $\lambda := \lambda(\hat{\zeta}) = e^{-\hat{\zeta}}$, $\bar{\lambda} := \bar{\lambda}(\bar{\zeta}) = e^{-\bar{\zeta}}$ and define $C := C(\zeta, k, K) = \{e^{\hat{\zeta}} \cdot C_f\}^{kK}$ where $C_f > 1$ be a bound on the norm of the derivative Df. Let $\epsilon_0^2 = \epsilon_0(\lambda, \bar{\lambda}, C, K)$ and $\bar{C}_2 = \bar{C}(\bar{\lambda}, C, K) > 0$ be the ϵ_0 and $\bar{C}$ as in Proposition 2.2 for this Δ.

Note that dominated splitting is always continuous (see [2]). So we can take $\gamma > 0$ small enough such that if for any $x, y \in \Delta$, if $\rho(x, y) < \gamma$, then for all $1 \le i \le K$,

$$e^{i(\hat{\zeta}-\zeta)} \le \frac{\|Df^i|_E(x)\|}{\|Df^i|_E(y)\|}, \ \frac{\|Df^i|_F(x)\|}{\|Df^i|_F(y)\|} \le e^{i(\zeta-\hat{\zeta})}. \tag{3.3}$$

Take $\sigma = \min\{\tau, \gamma\}$. If $z \in M$ and there exists pseudo-orbit $\{x_i,\, n_i\}_{i=-\infty}^{+\infty}$ satisfying $n_i \ge 2kK$ and $x_i, f^{n_i}(x_i) \in \Lambda_k(K, \zeta)(\forall\, i)$ such that z is a σ-shadowing point for $\{x_i, n_i\}_{i=-\infty}^{+\infty}$. $\rho\big(f^{c_i+j}(z), f^j(x_i)\big) \le \sigma \le \tau$ implies that the orbit of z is contained in the $\tau-$neighborhood U of Δ_1, and thus $z \in \Delta$. Since $x_i, f^{n_i}(x_i) \in \Lambda_k(K,\, \zeta)$ and $\rho\big(f^{c_i+j}(z), f^j(x_i)\big) \le \sigma \le \gamma$, it is easy to verify, using inequality (3.3), that z satisfies the condition as in Proposition 2.2 for the above λ and C. So by the choice of ϵ_0^2, the center stable and center unstable manifolds of all such points z above are the truth stable and unstable manifolds, and their sizes are at least ϵ_0^2.

Using Corollary 2.3 for the above set Δ, there exists $0 < \epsilon_0 \le \min\{\epsilon_0^1, \epsilon_0^2\}$ such that for any $0 < \epsilon \le \epsilon_0$, there exists $\delta = \delta(\epsilon)$ such that for any $x, y \in \Delta$, if $\rho(x, y) < \delta$, then $W_\epsilon^{cs}(x)$ and $W_\epsilon^{cu}(y)$ are transversal and have a unique intersection point. Clearly, $\bar{C}$ can be chosen $\max\{\bar{C}_1, \bar{C}_2\}$.

Since $\epsilon_0 \le \min\{\epsilon_0^1, \epsilon_0^2\}$, then for any $0 < \epsilon \le \epsilon_0$, the center stable and center unstable manifolds of all $x \in \Lambda_k^*(K, \zeta, \sigma)$ are truth stable and unstable manifold, and their sizes are at least ϵ. Note that $\Lambda_k^*(K, \zeta, \sigma) \subseteq \Delta$. Thus for any $x, y \in \Lambda_k^*(K, \zeta, \sigma) \subseteq \Delta$, if $\rho(x, y) < \delta$, then the stable manifold $W_\epsilon^{cs}(x)$ and unstable manifold $W_\epsilon^{cu}(y)$ are transversal and have a unique intersection point. □

Remark 3.5. The discussion of stable manifold for hyperbolic ergodic measures whose Oseledec splitting is dominated also appeared in [1]. But the result in [1] only gave the existence of stable manifold for a.e. points. Here in our paper we give the clear and definite block such that all points in the block have the same size of stable manifold and so do the (periodic) points nearby. These analysis will play crucial roles in the proof of Theorem 1.6. Furthermore, we point out that from the above proof, if σ is small enough, one can get a more strong relation: $\Lambda_k^*(K, \zeta, \sigma) \subseteq \Lambda(K', \zeta')$ for some small $\zeta' > 0$ and some large K'.

4 (Exponentially) Shadowing property

In this section we show exponentially shadowing property on Pesin blocks.

Proposition 4.1. *Assume* $\Lambda_k(K,\,\zeta)\neq\emptyset$ *for some* $k,\,K\in\mathbb{N}$ *and* $\zeta>0$*. Then* $\Lambda_k(K,\,\zeta)$ *satisfies (exponentially) shadowing property as the result (i) in Theorem 1.3(Replacing* Λ_τ *by* $\Lambda_k(K,\,\zeta)$ *and replacing* θ_τ *by some* $\theta>0$ *which is only dependent on* ζ*).*

Before proving that we need the shadowing lemma on nonempty Pesin blocks, which is obtained in [15].

Theorem 4.2. *(Shadowing lemma) If* $\Lambda_k(K,\,\zeta)\neq\emptyset$ *for some* $k,\,K\in\mathbb{N}$ *and* $\zeta>0$*, then* $\Lambda_k(K,\,\zeta)$ *satisfies shadowing property in following sense. For* $\forall\ \varepsilon>0$*, there exists* $\delta>0$ *such that if a* δ*-pseudo-orbit* $\{x_i,\,n_i\}_{i=-\infty}^{+\infty}$ *satisfies* $n_i\geq 2kK$ *and* $x_i, f^{n_i}(x_i)\in\Lambda_k(K,\,\zeta)$ *for all* i*, then there exists a* ε*-shadowing point* $x\in M$ *for* $\{x_i,n_i\}_{i=-\infty}^{+\infty}$*. If further* $\{x_i,n_i\}_{i=-\infty}^{+\infty}$ *is periodic, i.e., there exists an* $m>0$ *such that* $x_{i+m}=x_i$ *and* $n_{i+m}=n_i$ *for all* i*, then the shadowing point* x *can be chosen to be periodic.*

Proof Note that every Pesin block in present paper is contained in the original Pesin set defined in [15] and every original Pesin set satisfies shadowing property. So every Pesin block in present paper also satisfies shadowing property. □

Remark 4.3. The shadowing can be Lipschitz shadowing. That is, there exists $L_k>0,\delta_k>0$ such that for any $0<\delta<\delta_k$ if a δ-pseudo-orbit $\{x_i,\,n_i\}_{i=-\infty}^{+\infty}$ satisfies $n_i\geq 2kK$ and $x_i,f^{n_i}(x_i)\in\Lambda_k(K,\,\zeta)$ for all i, then there exists a $L_k\delta$-shadowing point $x\in M$ for $\{x_i,n_i\}_{i=-\infty}^{+\infty}$. The main observation is that the used technique is Liao's shadowing lemma[11, 4] for quasi-hyperbolic orbit segments and Liao's shadowing lemma is Lipschitz shadowing. So we can follow the proof of shadowing lemma in [15] to prove. Here we omit the details.

Using Theorem 4.2 and Proposition 3.4, we start to prove exponentially shadowing property.

Proof of Proposition 4.1 Let $\bar\zeta$, $\sigma=\sigma(K,\zeta,\bar\zeta,k)>0,\epsilon_0=\epsilon(K,\zeta,\bar\zeta,k)>0$ and $\bar C=\bar C(K,k,\zeta,\bar\zeta)>0$ be the numbers as in Proposition 3.4. Moreover, let Δ be the invariant set in the proof of Proposition 3.4, which contains $\Lambda_k^*(K,\zeta,\sigma)$.

For any fixed $\eta\in(0,\sigma)$, take $\epsilon=\epsilon(\eta)>0$ small enough such that

$$diam(W^{cs/cu}_{\epsilon}(y)<\frac{\eta}{2\bar C}$$

for all $y\in\Delta$. By Corollary 2.4, for this ϵ we can take $b_1=b_1(\epsilon)\in(0,\eta)$ such that for $x\in\Delta$, if $y\in W^{cu}_\epsilon(x)$ and $\rho(f(x),f(y))<b_1$ then $f(y)\in W^{cu}_\epsilon(f(x))$.

For b_1, take $\epsilon_*=\epsilon(b_1)>0$ small enough such that

$$diam(W^{cs/cu}_{\epsilon_*}(y)<\frac{b_1}{2\bar C}$$

for all $y \in \Delta$. Moreover, by Proposition 3.4, we can take $b_2 = b_2(\epsilon_*) \in (0, \min\{\frac{b_1}{2}, \sigma\})$ such that for any $x, y \in \Lambda_k^*(K, \zeta, \sigma)$ if $\rho(x, y) < b_2$, then $W_{\epsilon_*}^{cs}(x)$ and $W_{\epsilon_*}^{cu}(y)$ are transversal and have a unique intersection point.

By Theorem 4.2, for b_2 there exists $\delta = \delta(b_2) > 0$ such that if a δ-pseudo-orbit $\{x_i, n_i\}_{i=-\infty}^{+\infty}$ satisfies $n_i \geq 2kK$ and $x_i, f^{n_i}(x_i) \in \Lambda_k(K, \zeta)$ for all i, then there exists a b_2-shadowing point $x \in M$ for $\{x_i, n_i\}_{i=-\infty}^{+\infty}$. If further $\{x_i, n_i\}_{i=-\infty}^{+\infty}$ is periodic, i.e., there exists an $m > 0$ such that $x_{i+m} = x_i$ and $n_{i+m} = n_i$ for all i, then the shadowing point x can be chosen to be periodic. We only need to prove for orbit segment $\{x_0, n_0\}$, $\{x, n_0\}$ is exponentially shadowing $\{x_0, n_0\}$, since the others are similar. That $b_2 < \sigma$ implies $x \in \Lambda_k^*(K, \zeta, \sigma)$. Notice that $\rho(x_0, x) < b_2$ and $x_0 \in \Lambda_k(K, \zeta) \subseteq \Lambda_k^*(K, \zeta, \sigma)$. So $W_{\epsilon_*}^{cs}(x_0)$ and $W_{\epsilon_*}^{cu}(x)$ are transversal and have a unique intersection point $y \in M$. Since $x_0 \in \Lambda_k^*(K, \zeta, \sigma)$ and $y \in W_{\epsilon_*}^{cs}(x_0)$, by Proposition 3.4 we have

$$\rho(f^j(x_0), f^j(y)) \leq diam(f^n W_{\epsilon_*}^{cs}(x_0)) \leq \bar{C} e^{-\bar{\zeta} j} diam(W_{\epsilon_*}^{cs}(x_0)) \leq \frac{b_1}{2} e^{-\bar{\zeta} j},\ j = 0, 1, \cdots, n_0.$$

Combing this inequality with $\rho(f^j(x_0), f^j(x)) \leq b_2 < \frac{b_1}{2}$, $j = 0, 1, \cdots, n_0$, we have

$$\rho(f^j(x), f^j(y)) \leq \rho(f^j(x_0), f^j(x)) + \rho(f^j(x_0), f^j(y)) \leq b_1,\ j = 0, 1, \cdots, n_0.$$

From the choice of b_1 and $f^j(x) \in f^j(\Lambda_k^*(K, \zeta, \sigma)) \subseteq f^j(\Delta) = \Delta$, by induction it is easy to get $f^j(y) \in W_\epsilon^{cu}(f^j(x))\, j = 0, 1, \cdots, n_0$. Note that $f^{n_0}(x)$ is also in $\Lambda_k^*(K, \zeta, \sigma)$. Using $f^{n_0}(y) \in W_\epsilon^{cu}(f^{n_0}(x))$, by Proposition 3.4 one can get

$$\rho(f^j(x), f^j(y)) \leq diam(f^{(n_0-j)} W_\epsilon^{cu}(x_0)) \leq \bar{C} e^{-\bar{\zeta}(n_0-j)} diam(W_\epsilon^{cu}(x_0)) \leq \frac{\eta}{2} e^{-\bar{\zeta}(n_0-j)},$$

$j = 0, 1, \cdots, n_0$. So

$$\rho(f^j(x_0), f^j(x)) \leq \rho(f^j(x_0), f^j(y)) + \rho(f^j(x), f^j(y))$$

$$\leq (\frac{\eta}{2} + \frac{b_1}{2}) \max\{e^{-\bar{\zeta} j}, e^{-\bar{\zeta}(n_0-j)}\} \leq \eta \max\{e^{-\bar{\zeta} j}, e^{-\bar{\zeta}(n_0-j)}\},\ j = 0, 1, \cdots, n_0.$$

Take $\theta = \bar{\zeta}$ and we complete the proof. ☐

5 Proof of Theorem 1.3

In this section we show a theorem that there is Pesin set with measure arbitrarily close to 1 and after that we combine Proposition 3.4 and Proposition 4.1 to prove Theorem 1.3. Now we firstly state the theorem about the existence of arbitrarily large-measured(close to 1) Pesin set.

Theorem 5.1. *Let us make the same assumptions as in Theorem 1.3. Then*

$$\mu(\cup_{\zeta>0} \cup_{K\in\mathbb{N}} \Lambda(K, \zeta)) = 1.$$

If further all the Lyapunov exponents of μ are far from zero, then there exists $\zeta_0 > 0$ such that for all $\zeta \in (0, \zeta_0)$

$$\mu(\cup_{K\in\mathbb{N}}\Lambda(K,\,\zeta)) = 1.$$

Moreover, if μ is ergodic, then there exists $K_0 \in \mathbb{N}$, $\zeta_0 > 0$ such that for all $K \geq K_0$, $\zeta \in (0, \zeta_0)$, $\mu(\Lambda(K,\,\zeta)) = 1$.

To prove this theorem we need a lemma as follows.

Lemma 5.2. *Let $f \in \mathrm{Diff}^1(M)$ preserve an invariant probability measure μ, and $E \subseteq TM$ be a $Df-$invariant subbundle defined over an $f-$invariant set with μ full measure. Let $\lambda_E^+(x)$ be the maximal Lyapunov exponent in $E(x)$ of the measure μ. Then, for any $\tau > 0, \varepsilon > 0$ there exist an invariant set $B_{\tau,\varepsilon}$ with $\mu(B_{\tau,\varepsilon}) \geq 1 - \tau$ and an integer $K_{\tau,\varepsilon}$ such that for every point $x \in B_{\tau,\varepsilon}$ and any $K \geq K_{\tau,\varepsilon}$, the Birkhoff averages*

$$\frac{1}{lK}\sum_{j=0}^{l-1}\log\|Df^K|_{E(f^{jK}(x))}\|$$

converge towards a number contained in $[\lambda_E^+(x), \lambda_E^+(x) + \varepsilon)$, when $l \to +\infty$.

The similar property holds for the minimal Lyapunov exponent.

Remark. The ergodic case firstly appeared in [1] and and the ergodicity is useful in their proof. And similar results can be hold for continuous linear cocycles on homeomorphisms(even continuous systems).

Proof By Oseledec Theorem, one has

$$\lim_{l\to+\infty}\int|\tfrac{\log\|Df^l|_{E(x)}\|}{l} - \lambda_E^+(x)|d\mu = \int\lim_{l\to+\infty}|\tfrac{\log\|Df^l|_{E(x)}\|}{l} - \lambda_E^+(x)|d\mu = 0.$$

We can always assume that following limits exist since they always exist for μ a.e. For any $\varepsilon > 0$ and $\tau \in (0, 1)$, we can take large K_1 such that $\int h(x)d\mu < \frac{\tau}{4}\varepsilon$ where

$$h(x) := |\frac{\log\|Df^{K_1}|_{E^s(x)}\|}{K_1} - \lambda_E^+(x)|.$$

Since μ is f^{K_1} invariant, $\frac{1}{l}\sum_{i=0}^{l-1} h(f^{iK_1}(x))$ converge a.e. to integrable nonnegative functions $h^*(x)$. Also $h^*(f^{K_1}(x)) = h^*(x)$ a.e. and $\int h^*(x)d\mu = \int h(x)d\mu < \tau\varepsilon$. We set $B_1 = \{x \in M|\, h^*(x) < \varepsilon\}$ and claim $\mu(B_1) \geq 1 - \tau$. In fact, if $\mu(B_1) < 1 - \tau$, then $\int h^*(x)d\mu \geq \int_{M\setminus B_1} h^*(x)d\mu \geq \varepsilon\mu(M \setminus B_1) > \tau\varepsilon$, a contradiction to the inequality $\int h^*(x)d\mu \leq \tau\varepsilon$.

Let

$$B_{\tau,\varepsilon} := B_1 \cup f(B_1) \cup \cdots \cup f^{K_1}(B_1),$$

then $fB_{\tau,\varepsilon} = B_{\tau,\varepsilon}$ and $\mu(B_{\tau,\varepsilon}) \geq \mu(B_1) \geq 1 - \tau$. For $K \geq 1$ and $x \in B_{\tau,\varepsilon}$, one decomposes the orbit segment of length K of x as $(x, f(x), \cdots, f^{j-1}(x))$, $(f^j(x), \cdots, f^{j+(r-1)K_1-1}(x))$, and $(f^{j+(r-1)K_1}(x), \cdots, f^{N-1}(x))$ such that $0 \leq j < K_1$, $j + rK_1 \geq K$ and all the pints $f^i(x), f^{j+K_1}(x), \cdots, f^{j+rK_1}(x)$ belong to B_1. One deduces that

$$\|Df^K|_{E(x)}\| \leq$$

$\|Df^j|_{E(x)}\| \cdot (\|Df^{K_1}|_{E(f^j(x))}\| \cdots \|Df^{K_1}|_{E(f^{j+(r-2)K_1}(x))}\|) \cdot \|Df^{K-(j+(r-1)K_1)}|_{E(f^{j+(r-1)K_1}(x))}\|$. Hence, for $x \in B_{\tau,\varepsilon}$ one has

$$\log \|Df^K|_{E(x)}\| \leq 2K_1 \cdot C_f + \sum_{s=0}^{r-2} \log \|Df^{K_1}|_{E(f^{j+sK_1}(x))}\|$$

where $C_f = max(\log \|Df\|, \log \|Df^{-1}\|)$.

The point $f^j(x) \in B_1$ implies that the average

$$\frac{1}{kK_1} \sum_{l=0}^{k-1} \log \|Df^{K_1}|_{E(f^{j+lK_1}(x))}\|$$

converges to a number in $[\lambda_E^+(x), \lambda_E^+(x) + \frac{\varepsilon}{2})$. Hence

$$\lim_{k\to+\infty} \frac{1}{kK} \sum_{l=0}^{k-1} \log \|Df^K|_{E(f^{lK}(x))}\| \leq \frac{2K_1 \cdot C_f}{K} + \lim_{k\to+\infty} \frac{1}{kK_1} \sum_{l=0}^{k-1} \log \|Df^{K_1}|_{E(f^{j+lK_1}(x))}\|$$

Therefore, taking $K_{\tau,\varepsilon} = \frac{4K_1 \cdot C_f}{\varepsilon}$ and for all $K \geq K_{\tau,\varepsilon}$, one gets

$$\lim_{k\to+\infty} \frac{1}{kK} \sum_{l=0}^{k-1} \log \|Df^K|_{E(f^{lK}(x))}\| < \lambda_E^+(x) + \varepsilon.$$

On the other hand, by sub-multiplications of the norms, we have

$$\lim_{k\to+\infty} \frac{1}{kK} \sum_{l=0}^{k-1} \log \|Df^K|_{E(f^{\delta lK}(x))}\| \geq \lim_{n\to+\infty} \frac{1}{n} \log \|Df^n|_{E(x)}\| = \lambda_E^+(x).$$

So we complete the proof. $\square$

Now we start to prove Theorem 5.1.

Proof of Theorem 5.1 Let $\lambda_s(x), \lambda_u(x)$ denote the maximal and minimal Lyapunov exponents of $E^s(x)$ and $E^u(x)$ respectively for μ a.e. x. Notice that

$$\mu\big(\cup_{\zeta>0} \{x \in M \mid \lambda_s(x) < -3\zeta, \lambda_u(x) > 3\zeta\} \big) = 1$$

and

$$\mu\big(\cup_{\zeta>0} \cup_{S_0\geq 1} \{x \in M \mid \frac{1}{S} \log \frac{\|Df^S|_{E^s(y)}\|}{m(Df^S|_{E^u(y)})} \leq -2\zeta,\ \forall y \in Orb(x)\ S \geq S_0\} \big) = 1.$$

So for any $\tau \in (0,1)$, we can take small $\zeta > 0$, large S_0 and Δ_1 with $\mu(\Delta_1) > 1 - \frac{\tau}{2}$ such that if $x \in \Delta_1$ then $\lambda_s(x) < -3\zeta,\ \lambda_u(x) > 3\zeta$ and

$$\frac{1}{S} \log \frac{\|Df^S|_{E^s(y)}\|}{m(Df^S|_{E^u(y)})} \leq -2\zeta,\ \forall y \in Orb(x)\ S \geq S_0.$$

On the other hand, for $\varepsilon = \zeta$ in Lemma 5.2 we can take an invariant set $\Delta_2 = B_{\frac{\tau}{2},\varepsilon}$ with $\mu(\Delta_2) > 1 - \frac{\tau}{2}$ such that Δ_2 satisfies the result of Lemma 5.2 for both bundles $E^s(x)$ and $E^u(x)$. More precisely, there is $K_{\frac{\tau}{2},\varepsilon}$ such that for every point $x \in \Delta_2$ and any $K \geq K_{\frac{\tau}{2},\varepsilon}$, the Birkhoff averages satisfy

$$\lim_{l\to+\infty} \frac{1}{lK} \sum_{j=0}^{l-1} \log \|Df^K|_{E^s(f^{jK}(x))}\| \leq \lambda_s(x) + \varepsilon$$

and

$$\lim_{l\to+\infty} \frac{1}{lK} \sum_{j=0}^{l-1} \log m(Df^K|_{E^u(f^{jK}(x))}) \geq \lambda_u(x) - \varepsilon$$

Take $\Delta_\tau = \Delta_1 \cap \Delta_2$ and $K_\tau = \max\{S_0,\, K_{\frac{\tau}{2},\varepsilon}\}$. Clearly $\mu(\Delta_\tau) > 1 - \tau$. Then for every point $x \in \Delta_\tau$ and any $K \geq K_\tau$, we have

$$\lim_{l\to+\infty} \frac{1}{lK} \sum_{j=0}^{l-1} \log \|Df^K|_{E^s(f^{jK}(x))}\| \leq \lambda_s(x) + \varepsilon \leq -2\zeta,$$

$$\lim_{l\to+\infty} \frac{1}{lK} \sum_{j=0}^{l-1} \log m(Df^K|_{E^u(f^{jK}(x))}) \geq \lambda_u(x) - \varepsilon \geq 2\zeta$$

and

$$\frac{1}{L} \log \frac{\|Df^L|_{E(y)}\|}{m(Df^L|_{F(y)})} \leq -2\zeta,\ \forall y \in Orb(x),\ L \geq K.$$

Obviously the last inequality is same as condition (c) in the definition of Pesin set $\Lambda(K,\zeta)$. Since conditions (a) and (b) in the definition of Pesin set $\Lambda(K,\zeta)$ are same as in [15], using the same discussion as in [15] we can get

$$\Delta_\tau \subseteq \bigcup_{k\geq 1} \Lambda_k(K,\zeta)$$

and then $\Delta_\tau \subseteq \Lambda(K,\zeta)$(Note that both Δ_τ and $\Lambda(K,\zeta)$ are invariant). This process is easy, only based on the definition of the above two limit inequalities, and here omit the details.

By the arbitrary choice of small τ we complete the proof. $\square$

At the end of this section we prove Theorem 1.3.

Proof of Theorem 1.3 Fix $\tau > 0$. By Theorem 5.1 we can take $\zeta > 0$ small enough and K large enough such that $\mu(\Lambda(K,\, \zeta)) > 1 - \tau$. Let

$$\Lambda_\tau = \Lambda(K,\, \zeta).$$

Then by Proposition 3.4 and Proposition 4.1, take $\theta_\tau = \theta(\zeta) > 0,\ T_\tau = 2kK \in \mathbb{N}$ and thus Λ_τ is the needed set. $\square$

6 Proof of Theorem 1.6 and 1.8

In this section we use Theorem 1.5 to prove Theorem 1.6 and 1.8. Note that different to Katok's (shadowing and) closing lemma, the length of every pseud-orbit segment in the present paper must be larger than T_τ and thus we must take care when we use (shadowing and) closing lemma. Firstly, we prove Theorem 1.6.

Proof of Theorem 1.6 The proof is an adaption of [16]. Since hyperbolic measure can be always divided into at most $D = dim(M)$ hyperbolic measures with fixed indexs, we can assume μ is hyperbolic measure with fixed index.

Let $x \in supp(\mu)$ and let $\eta > 0$ be small enough. There exists a positive number τ small enough such that $\mu(B(x, \eta/2) \cap \Lambda_\tau) > 0$. By Theorem 1.5, we can take $\sigma > 0$ such that the stable and unstable manifolds of $\Lambda_\tau^*(\sigma)$ defined in Theorem 1.5 have the same uniform size. Let $\delta > 0$ be the number such that if $x, y \in \Lambda_\tau^*(\sigma)$ are δ close, then their stable and unstable manifolds are transversal. Let T_τ be the number as in Theorem 1.5 which is only dependent on Λ_τ^*. Now let $r > 0$ be an arbitrary small number, with $r < \min\{\eta/2, \sigma, \delta/3\}$.

Pick a set $B \subseteq B(x, \eta/2) \cap \Lambda_\tau$ of diameter less than $\beta = \beta(\tau, r)$ (as in Theorem 1.5) less than r and of positive measure. Let $x_1 \in B$ be a recurrent point (by the Poincare Recurrence Theorem), and $n(x_1) \geq T_\tau$ a positive integer such that $f^{n(x_1)}(x_1) \in B$. Since $\rho(x_1, f^{n(x_1)}(x_1)) < \beta$, in applying Theorem 1.5 we obtain that there exists a periodic point z_1 of period $n(x_1)$ such that $\rho(f^i(x_1), f^i(z_1)) < r < \min\{\eta/2, \sigma\}$, $0 \leq i \leq n(x_1) - 1$. This implies that $z_1 \in \Lambda_\tau^*(\sigma)$ and $\rho(x, z_1) < \eta$.

We will prove that μ-almost every point of B belongs to $\overline{W^u(z_1)}$. For that, let $y \in B$ be a Borel density point, and let $\tau' > 0$ be small enough such that $\tau' < r/2$, $\rho(z_1, y) > \tau'$ and $\mu(B \cap B(y, \tau'/2)) > 0$. Pick a set $\tilde{B} \subseteq B \cap B(y, \tau'/2)$ of diameter less than $\tilde{\beta} = \beta(k, \tau'/2)$ (as in Theorem 1.5) and of positive measure. Let $x_2 \in \tilde{B}$ be a recurrent point, and $n(x_2) \geq T_\tau$ a positive integer such that $f^{n(x_2)}(x_2) \in B$. Since $\rho(x_2, f^{n(x_2)}(x_2)) < \tilde{\beta}$. Since $\rho(x_2, f^{n(x_2)}(x_2)) < \tilde{\beta}$, in applying Theorem 1.5 we obtain that there exists a periodic point z_2 of period $n(x_2)$ such that $\rho(f^i(x_2), f^i(z_2)) < \tau'/2 < r < \sigma$, $0 \leq i \leq n(x_2) - 1$. This implies that $z_2 \in \Lambda_\tau^*(\sigma)$ and $\rho(y, z_2) < \rho(y, x_2) + \rho(x_2, z_2) < \tau'$, thus $z_1 \neq z_2$.

Note that

$$\rho(z_1, z_2) < \rho(x_1, z_1) + \rho(x_1, y) + \rho(y, z_2) < r + r + \tau' < 3r < \delta.$$

By the choice of δ, the stable (unstable) manifold $W^{s/u}(z_1)$ and unstable (stable) manifold $W^{u/s}(z_2)$ are transversal. So z_1 has homoclinic point and hence there exists horseshoe.

Let w be a transverse intersection point on $W^u(z_1)$ and $W^s(z_2)$. By the Inclination Lemma(see e.g. [9]), we have that $W^u(z_1)$ accumulates locally on $W^u(z_2)$. Indeed, let N be a common period for z_1 and z_2. The points z_1 and z_2 are hyperbolic fixed points for f^N with w a transverse intersection point of $W^u(z_1)$ and $W^s(z_2)$. The images under f^N of a ball around w in $W^u(z_1) = f^N W^u(z_1)$ accumulate on $W^u(z_2)$. Hence

$$\rho(W^u(z_1), y) \leq \rho(W^u(z_1), x_2) + \rho(x_2, y) \leq \tau'$$

for all $\tau' > 0$. This implies that $y \in \overline{W^u(z_1)}$, for all Borel density points $y \in B$. Therefore $\mu(\overline{W^u(z_1)}) \geq \mu(B) > 0$.

Since the hyperbolic periodic point z_1 is such that $\rho(x, z_1) < \eta,$, where x is an arbitrary point of $supp(\mu)$ and $\eta > 0$ is arbitrarily small, we obtain that the support of the measure μ is contained in the closure of all hyperbolic periodic points with $\mu(\overline{W^u(z_1)}) > 0$. $\square$

Secondly, we start to prove Theorem 1.8.

Proof of Theorem 1.8 The proof is an adaption of [8, 10, 3] and here we only prove the ergodic case similar as Theorem 15.3.1 in [3]. Then there is $x \in supp(\mu)$ such that $supp(\mu) = \overline{orb(x)}$(In fact this property holds for μ a.e. x, see Theorem 5.15 in [17]). Given $\tau > 0$ and let Λ_τ be as in Theorem 1.5. We may assume that $supp(\mu) = \overline{orb(x)}$ for some $x \in \Lambda_\tau$ and that for every $\tau > 0$, the intersection $orb(x) \cap \Lambda_\tau$ is dense in Λ_τ. Let $T_\tau > 0$ is the number only dependent on τ in Theorem 1.5. Set $\psi = 0$ and

$$\psi(f^n(x)) = \sum_{k=0}^{n-1} \varphi(f^k(x))$$

for each $n \in \mathbb{N}$. To extend ψ continuously to Λ_τ we show that ψ is uniformly continuous on $orb(x) \cap \Lambda_\tau$. For any $\eta > 0$, let $\beta = \beta(\tau, \eta) > 0$ be as in Theorem 1.5.

Firstly, we consider $n_2 > n_1$ satisfying $n_2 - n_1 \geq T_\tau$, $f^{n_1}(x), f^{n_2}(x) \in \Lambda_\tau$ and $\rho(f^{n_1}(x),\ f^{n_2}(x)) < \beta$, then by Theorem 1.5, there exists a hyperbolic periodic point z with period $n_2 - n_1$ such that

$$\rho(f^j(f^{n_1}(x)),\ f^j(z)) < \eta \cdot e^{-\min\{j, n_2 - n_1 - j\}\theta_\tau},$$

where $j = 0, 1, \cdots, n_2 - n_1 - 1$ and $\theta_\tau > 0$ is the number only dependent on τ in Theorem 1.5. Since φ is Hölder continuous,

$$|\varphi(x) - \varphi(y)| \leq C\rho(x, y)^\kappa$$

for some $C > 0$ and $0 < \kappa \leq 1$, and we obtain

$$|\varphi(f^j(f^{n_1}(x)) - \varphi(f^j(z))| \leq C\eta^\kappa \cdot e^{-\kappa \min\{j, n_2 - n_1 - j\}\theta_\tau}.$$

Thus,

$$\begin{aligned}
& |\psi(f^{n_1}(x)) - \psi(f^{n_2}(x))| \\
= \ & |\sum_{j=n_1}^{n_2-n_1-1} (\varphi(f^j(f^{n_1}(x)) - \varphi(f^j(z))) + \sum_{j=n_1}^{n_2-n_1-1} \varphi(f^j(z))| \\
\leq \ & C\eta^\kappa \cdot \sum_{j=n_1}^{n_2-n_1-1} \max\{e^{-j\kappa\theta_\tau}, e^{-(n_2-n_1-j)\kappa\theta_\tau}\} + |\sum_{j=n_1}^{n_2-n_1-1} \varphi(f^j(z))| \\
\leq \ & C\eta^\kappa N
\end{aligned}$$

for some constant $N > 0$ independent of n_1 and n_2.

Secondly, we consider $n_2 > n_1$ satisfying $f^{n_1}(x), f^{n_2}(x) \in \Lambda_\tau$ and $\rho(f^{n_1}(x),\ f^{n_2}(x)) < \frac{\beta}{2}$. Since $orb(x) \cap \Lambda_\tau$ is dense in Λ_τ, then we can take n_3 large enough such that $n_3 -$

$n_1 > n_3 - n_2 \geq T_\tau$, $f^{n_3}(x) \in \Lambda_\tau$ and $\rho(f^{n_3}(x),\ f^{n_1}(x)) < \frac{\beta}{2}$. So $\rho(f^{n_3}(x),\ f^{n_2}(x)) \leq \rho(f^{n_3}(x),\ f^{n_1}(x)) + \rho(f^{n_1}(x), f^{n_2}(x)) < \beta$. Thus by the above discussion, we have

$$|\psi(f^{n_3}(x)) - \psi(f^{n_1}(x))| \leq C\eta^\kappa N$$

and

$$|\psi(f^{n_3}(x)) - \psi(f^{n_2}(x))| \leq C\eta^\kappa N.$$

Therefore,

$$\begin{aligned} & |\psi(f^{n_1}(x)) - \psi(f^{n_2}(x))| \\ \leq\ & |\psi(f^{n_3}(x)) - \psi(f^{n_1}(x))| + |\psi(f^{n_3}(x)) - \psi(f^{n_2}(x))| \\ \leq\ & 2C\eta^\kappa N. \end{aligned}$$

This shows that we can extend ψ continuously to Λ_τ. Now we extend ψ to $\cup_{i=0}^{\infty} f^i(\Lambda_\tau)$ as follows. If $y \in f(\Lambda_\tau) \setminus \Lambda_\tau$, then let

$$\psi(y) = \psi(f^{-1}(y)) + \varphi(f^{-1}(y)),$$

and we use the same expression for all $y \in \cup_{i=0}^{n+1} f^i(\Lambda_\tau) \setminus \cup_{i=0}^{n} f^i(\Lambda_\tau)$. Since $\mu(\cup_{i=0}^{\infty} f^i(\Lambda_\tau)) = 1$, the function ψ is defined almost everywhere and clearly satisfies the needed condition. $\square$

We point out that the stable manifold is useless in the proof in Theorem 1.8 and exponentially closing plays the crucial role.

References.

1. F. Abdenur, C. Bonatti, S. Crovisier, *Nonuniform hyperbolicity of C^1-generic diffeomorphisms*, Israel Journal of Mathematics, to appear.
2. Bonatti, Lorenzo, Viana, *Dynamics beyond uniform hyperbolicity: a global geometric and probabilistic perspective*, Springer-Verlag Berlin Heidelberg (2005), 287-293.
3. Luis Barreira, Yakov B. Pesin, *Nonuniform hyperbolicity*,Cambridge Univ. Press, Cambridge (2007).
4. S. Gan, *A generalized shadowing lemma*, Dist. Cont. Dyn. Sys. 8 (2002), 627-632.
5. S. Gan, *Horseshoe and entropy for C^1 surface diffeomorphisms*, Nonlinearity 15 (2002) 841-848.
6. M. Hirsch, C. Pugh and M. Shub, *Invariant Manifolds* (Lecture Notes in Math. vol 583)(Berlin: Spinger),1977.
7. A. Katok, *Liapunov exponents, entropy and periodic orbits for diffeomorphisms*, Pub. Math. IHES, 51 (1980), 137-173.
8. A. Katok, *Nonuniform hyperbolicity an stucture of smooth dynamical systems*, Proceedings of the International Congress of Mathematicians (Warsaw, 1983), PWN, 1984, 1245-1253.

9. A. Katok and B. Hasselblatt, *Introduction to the modern theory of dynamical systems*, Encyclopedia of Mathematics and its Applications 54, Cambridge Univ. Press, Cambridge (1995).
10. A. Katok and L. Mondoz, *Dynamical systems with nonuniformly hyperbolic behavior*, to supplement to *Introduction to the modern theory of dynamical systems,* by A. Katok and B. Hasselblatt, Cambridge Univ. Press, Cambridge (1995).
11. S. Liao, *An existence theorem for periodic orbits*, Acta Sci. Natur. Univ. Pekin. (1979), 1-20.
12. V. I. Oseledec, *Multiplicative ergodic theorem, Liapunov characteristic numbers for dynamical systems*, Trans. Moscow Math. Soc., 19 (1968), 197-221; translated from Russian.
13. M. Pollicott, *Lectures on ergodic theory and Pesin theory on compact manifolds*, Cambridge Univ. Press, Cambridge (1993).
14. Pujals E and Sambarino M, *Homoclinic tangencies and hyperbolicity for surface diffeomorphisms: a conjecture of Palis*, Ann. Math. 151 (2000), 961-1023
15. W. Sun, X. Tian, *Pesin set, closing lemma and shadowing lemma in C^1 non-uniformly hyperbolic systems with limit domination*, arXiv:1004.0486.
16. I. Ugarcovici, *On hyperbolic measures and periodic orbits*, Discrete and continuous dynamical systems, 16(2006), 505-512.
17. P. Walters, *An introduction to ergodic theory*, Springer-Verlag, 2001.
18. J. Yang, *C^1 dynamics far from tangencies*, preprint.